arXiv:math/0502270v1 [math.GR] 13 Feb 2005

A CLASS OF RIGID COXETER GROUPS

TETSUYA HOSAKA

Abstract. In this paper, we give a new class of rigid Coxeter groups. Let (W, S) be a Coxeter system. Suppose that (0) for each $s, t \in S$ such that $m(s, t)$ is even, $m(s, t) = 2$, (1) for each $s \neq t \in S$ such that $m(s, t)$ is odd, $\{s, t\}$ is a maximal spherical subset of S, (2) there does not exist a three-points subset $\{s, t, u\} \subset S$ such that $m(s, t)$ and $m(t, u)$ are odd, and (3) for each $s \neq t \in S$ such that $m(s, t)$ is odd, the number of maximal spherical subsets of S intersecting with $\{s, t\}$ is at most two, where $m(s, t)$ is the order of st in the Coxeter group W. Then we show that the Coxeter group W is rigid. This is an extension of a result of D. Radcliffe.

1. Introduction and preliminaries

The purpose of this paper is to give a new class of rigid Coxeter groups. A *Coxeter group* is a group W having a presentation

$$\langle\, S \,|\, (st)^{m(s,t)} = 1 \text{ for } s, t \in S \,\rangle,$$

where S is a finite set and $m : S \times S \to \mathbb{N} \cup \{\infty\}$ is a function satisfying the following conditions: (i) $m(s, t) = m(t, s)$ for any $s, t \in S$, (ii) $m(s, s) = 1$ for any $s \in S$, and (iii) $m(s, t) \geq 2$ for any $s, t \in S$ such that $s \neq t$. The pair (W, S) is called a *Coxeter system*. For a Coxeter group W, a generating set S' of W is called a *Coxeter generating set for* W if (W, S') is a Coxeter system. In a Coxeter system (W, S), the conjugates of elements of S are called *reflections*. Let (W, S) be a Coxeter system. For a subset $T \subset S$, W_T is defined as the subgroup of W generated by T, and called a *parabolic subgroup*. If T is the empty set, then W_T is the trivial group. A subset $T \subset S$ is called a *spherical subset of* S, if the parabolic subgroup W_T is finite.

A *diagram* is an undirected graph Γ without loops or multiple edges with a map $\mathrm{Edges}(\Gamma) \to \{2, 3, 4, \ldots\}$ which assigns an integer greater

---

Date: November 5, 2004.

2000 *Mathematics Subject Classification.* 20F65, 20F55.

Key words and phrases. rigidity of Coxeter groups.

Partly supported by the Grant-in-Aid for Scientific Research, The Ministry of Education, Culture, Sports, Science and Technology, Japan, (No. 15740029).

than 1 to each of its edges. Since such diagrams are used to define Coxeter systems, they are called *Coxeter diagrams*.

Let (W, S) and (W', S') be Coxeter systems. Two Coxeter systems (W, S) and (W', S') are said to be *isomorphic*, if there exists a bijection $\psi : S \to S'$ such that

$$m(s,t) = m'(\psi(s), \psi(t))$$

for every $s, t \in S$, where $m(s,t)$ and $m'(s',t')$ are the orders of st in W and $s't'$ in W', respectively.

In general, a Coxeter group does not always determine its Coxeter system up to isomorphism. Indeed some counter-examples are known.

Example 1 ([1, p.38 Exercise 8], [2])**.** It is known that for an odd number $k \geq 3$, the Coxeter groups defined by the diagrams in Figure 1 are isomorphic and D_{2k}.



FIGURE 1. Two distinct Coxeter diagrams for D_{2k}

Example 2 ([2])**.** It is known that the Coxeter groups defined by the diagrams in Figure 2 are isomorphic by the *diagram twisting* ([2, Definition 4.4]).



FIGURE 2. Coxeter diagrams for isomorphic Coxeter groups

Here there exists the following natural problem.

Problem ([2], [3])**.** When does a Coxeter group determine its Coxeter system up to isomorphism?

A Coxeter group W is said to be *rigid*, if for each Coxeter generating sets S and S' for W the Coxeter systems (W,S) and (W,S') are isomorphic.

The following theorem was proved by D. Radcliffe in [5].

Theorem 1.1 ([5], [6]). *If (W,S) is a right-angled Coxeter system (i.e. $m(s,t)=2$ for each $s \neq t \in S$), then the Coxeter group W is rigid.*

In this paper, we prove the following theorem which is an extension of Theorem 1.1.

Theorem 1.2. *Let (W,S) be a Coxeter system. Suppose that*

(0) *for each $s,t \in S$ such that $m(s,t)$ is even, $m(s,t)=2$,*
(1) *for each $s \neq t \in S$ such that $m(s,t)$ is odd, $\{s,t\}$ is a maximal spherical subset of S,*
(2) *there does not exist a three-points subset $\{s,t,u\} \subset S$ such that $m(s,t)$ and $m(t,u)$ are odd, and*
(3) *for each $s \neq t \in S$ such that $m(s,t)$ is odd, the number of maximal spherical subsets of S intersecting with $\{s,t\}$ is at most two.*

Then the Coxeter group W is rigid.

Here we can not omit the condition (3) in this theorem by Example 2.

Example 3. The Coxeter groups defined by the diagrams in Figure 3 are rigid by Theorem 1.2.



Figure 3. Coxeter diagrams for rigid Coxeter groups

2. Lemmas on Coxeter groups

We first recall some basic results about Coxeter groups.

Lemma 2.1 ([1]). *Let (W,S) be a Coxeter system and let $T \subset S$. If there exists $w \in W$ such that $\ell(tw) < \ell(w)$ for every $t \in T$, then T is a spherical subset of S. Here $\ell(w)$ is the length of w with respect to S.*

Lemma 2.2 ([1, p.12 Proposition 3]). *Let (W,S) be a Coxeter system and let $s,t \in S$. Then s is conjugate to t if and only if there exists a sequence $s_1,\dots,s_n \in S$ such that $s_1=s$, $s_n=t$ and $m(s_i,s_{i+1})$ is odd for each $i \in \{1,\dots,n-1\}$.*

Lemma 2.3 (cf. [2]). *Let (W, S) and (W', S') be Coxeter systems. Suppose that there exists an isomorphism $\phi : W \to W'$. Then for each maximal spherical subset $T \subset S$, there exists a unique maximal spherical subset $T' \subset S'$ such that $\phi(W_T) = w' W'_{T'} (w')^{-1}$ for some $w' \in W'$.*

Let (W, S) be a Coxeter system and let $w \in W$. A representation $w = s_1 \cdots s_l$ $(s_i \in S)$ is said to be *reduced*, if $\ell(w) = l$.

Using the above lemmas, we prove some lemmas needed later.

Lemma 2.4. *Let (W, S) be a Coxeter system which satisfies the conditions (0) and (1) in Theorem 1.2, let $w, v, x \in W$ and let $w = s_1 \cdots s_l$ and $v = t_1 \cdots t_m$ $(s_i, t_j \in S)$ be reduced representations. Suppose that $W_{\{s_1,\dots,s_l\}} \cong \mathbb{Z}_2^l$, $W_{\{t_1,\dots,t_m\}} \cong \mathbb{Z}_2^m$ and $l \geq 2$. If $w = xvx^{-1}$ then $w = v$.*

Proof. We prove this lemma by the induction on the length of x. If $\ell(x) = 0$ then $x = 1$ and we obtain that if $w = xvx^{-1}$ then $w = v$. Now we suppose that if $w = xvx^{-1}$ then $w = v$ for each $x \in W$ such that $\ell(x) < k$. Let $x \in W$ such that $\ell(x) = k$. Suppose that $w = xvx^{-1}$. Let $x = a_1 \cdots a_k$ $(a_i \in S)$ be a reduced representation. Then

$$s_1 \cdots s_l = (a_1 \cdots a_k)(t_1 \cdots t_m)(a_k \cdots a_1).$$

If $(a_1 \cdots a_k)(t_1 \cdots t_m)$ is not reduced, then we may suppose that $a_k = t_1$, and

$$(a_1 \cdots a_k)(t_1 \cdots t_m)(a_k \cdots a_1) = (a_1 \cdots a_{k-1})(t_1 \cdots t_m)(a_{k-1} \cdots a_1),$$

because $t_i t_j = t_j t_i$ for each i, j. Then by the inductive hypothesis, we obtain $w = v$. By the same argument, if $(s_1 \cdots s_l)(a_1 \cdots a_k)$ is not reduced, then we obtain $w = v$ by the inductive hypothesis. We suppose that $(a_1 \cdots a_k)(t_1 \cdots t_m)$ and $(s_1 \cdots s_l)(a_1 \cdots a_k)$ are reduced. Here

$$(s_1 \cdots s_l)(a_1 \cdots a_k) = (a_1 \cdots a_k)(t_1 \cdots t_m).$$

This implies that $l = m$. Let $z = (s_1 \cdots s_l)(a_1 \cdots a_k)$. Then $\ell(a_1 z) < \ell(z)$ and $\ell(s_i z) < \ell(z)$ for each $i = 1, \dots, l$. By Lemma 2.1, $\{s_1, \dots, s_l, a_1\}$ is a spherical subset of S. Since $l \geq 2$, $W_{\{s_1,\dots,s_l,a_1\}} \cong \mathbb{Z}_2^{l+1}$ by (0) and (1). Hence $a_1 s_i = s_i a_1$ for each $i = 1, \dots, l$ and

$$a_1 z = (s_1 \cdots s_l)(a_2 \cdots a_k) = (a_2 \cdots a_k)(t_1 \cdots t_m),$$

i.e., $w = (a_2 \cdots a_k) v (a_k \cdots a_2)$. By the inductive hypothesis, we obtain $w = v$. $\square$

Lemma 2.5. *Let (W, S) be a Coxeter system which satisfies the conditions (0), (1), (2) and (3) in Theorem 1.2, let $x \in W$ and let T and U*

be subsets of S such that $W_T \cong \mathbb{Z}_2^{|T|}$, $W_U \cong \mathbb{Z}_2^{|U|}$, $|T| \geq 2$ and $|U| \geq 2$. Then $W_T \cap xW_Ux^{-1} \subset W_{T\cap U}$.

Proof. Let $w \in W_T \cap xW_Ux^{-1}$. Then $w = xvx^{-1}$ for some $v \in W_U$. If $\ell(w) \geq 2$ then by Lemma 2.4, $w = v \in W_T \cap W_U = W_{T\cap U}$. We suppose that $\ell(w) = 1$, i.e., $w \in T$. Let $t = w$. Then $t = xvx^{-1}$ is a reflection and $v \in W_U$. Hence $t = xvx^{-1} = ysy^{-1}$ for some $s \in U$ and $y \in W$, i.e., $t \sim s$. Suppose that $t \neq s$. Then $m(t,s)$ is odd by Lemma 2.2 and (2). There exist maximal spherical subsets $\bar{T}$ and $\bar{U}$ of S such that $T \subset \bar{T}$ and $U \subset \bar{U}$. Here we note that $|\bar{T}| \geq |T| \geq 2$, $|\bar{U}| \geq |U| \geq 2$ and $\bar{T} \cap \bar{U} = \emptyset$ by (1). Then $\{t,s\}$, $\bar{T}$ and $\bar{U}$ are different maximal spherical subsets intersecting with $\{t,s\}$. This contradicts (3). Thus $t = s$ and $w = t = s \in T \cap U \subset W_{T\cap U}$. Hence $W_T \cap xW_Ux^{-1} \subset W_{T\cap U}$. □

Lemma 2.6. *Let (W,S) and (W',S') be Coxeter systems. Suppose that there exists an isomorphism $\phi : W \to W'$ and that*

- (2′) *there does not exist a three-points subset $\{s',t',u'\} \subset S'$ such that $m'(s',t')$ and $m'(t',u')$ are odd.*

Let $\{s,t\}$ and T be maximal spherical subsets of S such that $m(s,t)$ is odd. By Lemma 2.3, there exist maximal spherical subsets $\{s',t'\}$ and T' of S' such that $\phi(W_{\{s,t\}}) \sim W'_{\{s',t'\}}$ and $\phi(W_T) \sim W'_{T'}$. If $\{s,t\} \cap T \neq \emptyset$ then $\{s',t'\} \cap T' \neq \emptyset$.

Proof. Suppose that $\{s,t\} \cap T \neq \emptyset$. We may suppose that $t \in \{s,t\} \cap T$. Then $\phi(t) \in \phi(W_{\{s,t\}}) \sim W'_{\{s',t'\}}$. Since $m(s,t) = m'(s',t')$ is odd and $o(\phi(t)) = o(t) = 2$, $\phi(t)$ is a reflection in W' and $\phi(t) \sim s' \sim t'$. Here $\phi(t) = x't'x'^{-1}$ for some $x' \in W'$ and $\phi(W_T) = w'W'_{T'}w'^{-1}$ for some $w' \in W'$. Then

$$x't'x'^{-1} = \phi(t) \in \phi(W_T) = w'W'_{T'}w'^{-1}.$$

Hence $(w'^{-1}x')t'(x'^{-1}w') \in W'_{T'}$. We note that $(w'^{-1}x')t'(x'^{-1}w')$ is a reflection in $W'_{T'}$. Thus there exists $u' \in T'$ such that $t' \sim u'$. Since $s' \sim t' \sim u'$, we have that $u' \in \{s',t'\}$ by the hypothesis (2′) and Lemma 2.2. Hence $\{s',t'\} \cap T' \neq \emptyset$. □

3. Proof of the theorem

Let (W,S) be a Coxeter system. Suppose that

- (0) for each $s,t \in S$ such that $m(s,t)$ is even, $m(s,t) = 2$,
- (1) for each $s \neq t \in S$ such that $m(s,t)$ is odd, $\{s,t\}$ is a maximal spherical subset of S,

(2) there does not exist a three-points subset $\{s, t, u\} \subset S$ such that $m(s, t)$ and $m(t, u)$ are odd, and

(3) for each $s \neq t \in S$ such that $m(s, t)$ is odd, the number of maximal spherical subsets of S intersecting with $\{s, t\}$ is at most two.

Let (W', S') be a Coxeter system. We suppose that there exists an isomorphism $\phi : W \to W'$. The purpose is to show that the Coxeter systems (W, S) and (W', S') are isomorphic.

Lemma 3.1. *The Coxeter system (W', S') satisfies the following:*

($0'$) *for each $s', t' \in S'$ such that $m'(s', t')$ is even, $m'(s', t') = 2$,*

($1'$) *for each $s' \neq t' \in S'$ such that $m'(s', t')$ is odd, $\{s', t'\}$ is a maximal spherical subset of S',*

($2'$) *there does not exist a three-points subset $\{s', t', u'\} \subset S'$ such that $m'(s', t')$ and $m'(t', u')$ are odd, and*

($3'$) *for each $s' \neq t' \in S'$ such that $m'(s', t')$ is odd, the number of maximal spherical subsets of S' intersecting with $\{s', t'\}$ is at most two.*

Proof. ($0'$) and ($1'$) Let $s' \neq t' \in S'$. There exists a maximal spherical subset T' of S' such that $\{s', t'\} \subset T'$. By Lemma 2.3, $\phi^{-1}(W'_{T'}) \sim W_T$ for some maximal spherical subset T of S. By (0) and (1), either

(i) $W_T \cong \mathbb{Z}_2^{|T|}$, or

(ii) $|T| = 2$ and if $T = \{s, t\}$ then $m(s, t)$ is odd.

Hence W_T is a rigid Coxeter group and (W_T, T) and $(W'_{T'}, T')$ are isomorphic. Thus if $m'(s', t')$ is even then $m'(s', t') = 2$, and if $m'(s', t')$ is odd then $\{s', t'\}$ is a maximal spherical subset of S'. Hence ($0'$) and ($1'$) hold.

($2'$) Suppose that there exists a three-points subset $\{s', t', u'\} \subset S'$ such that $m'(s', t')$ and $m'(t', u')$ are odd. Then $\{s', t'\}$ and $\{t', u'\}$ are maximal spherical subsets of S' by ($1'$). By Lemma 2.3, there exist $s, t, u, v \in S$ such that $\phi^{-1}(W'_{\{s', t'\}}) \sim W_{\{s,t\}}$ and $\phi^{-1}(W'_{\{t', u'\}}) \sim W_{\{u,v\}}$, where $\{s, t\} \neq \{u, v\}$ (i.e., $|\{s, t, u, v\}| \geq 3$). We note that for each $x' \in W'_{\{s', t'\}}$, x' is a reflection on (W', S') if and only if $o(x') = 2$, because $m'(s', t')$ is odd. Hence for each $x' \in W'_{\{s', t'\}}$, x' is a reflection on (W', S') if and only if $\phi^{-1}(x')$ is a reflection on (W, S). Thus

$$s \sim t \sim \phi^{-1}(t') \sim \phi^{-1}(u') \sim u \sim v.$$

This contradicts (2) by Lemma 2.2, since $|\{s, t, u, v\}| \geq 3$. Thus ($2'$) holds.

($3'$) Let $\{s,t\}$ and T be maximal spherical subsets of S such that $m(s,t)$ is odd. By Lemma 2.3, there exist maximal spherical subsets $\{s',t'\}$ and T' of S' such that $\phi(W_{\{s,t\}}) \sim W'_{\{s',t'\}}$ and $\phi(W_T) \sim W'_{T'}$. Then by Lemma 2.6, (2) and ($2'$), $\{s,t\} \cap T \neq \emptyset$ if and only if $\{s',t'\} \cap T' \neq \emptyset$. Hence ($3'$) holds by (3). $\square$

Lemma 3.2. *Let* $T_1, \dots, T_k$ *be maximal spherical subsets of* S *and let* $T'_1, \dots, T'_k$ *be maximal spherical subsets of* S' *such that* $\phi(W_{T_i}) \sim W'_{T'_i}$ *and* $W_{T_i} \cong \mathbb{Z}_2^{|T_i|}$ *for each* i*. Then* $|T_1 \cap \cdots \cap T_k| = |T'_1 \cap \cdots \cap T'_k|$.

Proof. If $|T_{i_0}| = 1$ for some i_0, then $|T_1 \cap \cdots \cap T_k| = 0 = |T'_1 \cap \cdots \cap T'_k|$ because T_i and T'_i are maximal spherical subsets. We suppose that $|T_i| \geq 2$ for each i. Now $\phi(W_{T_i}) = w'_i W'_{T'_i} {w'}_i^{-1}$ for some $w'_i \in W'$, and

$$\phi(W_{T_1 \cap \cdots \cap T_k}) = (w'_1 W'_{T'_1} {w'}_1^{-1}) \cap \cdots \cap (w'_k W'_{T'_k} {w'}_k^{-1}) \subset w'_1 W'_{T'_1 \cap \cdots \cap T'_k} {w'}_1^{-1}$$

by Lemmas 2.5 and 3.1. Hence $\phi(W_{T_1 \cap \cdots \cap T_k}) \subset w'_1 W'_{T'_1 \cap \cdots \cap T'_k} {w'}_1^{-1}$. By applying the same argument to $\phi^{-1} : W' \to W$,

$$\phi^{-1}(W'_{T'_1 \cap \cdots \cap T'_k}) \subset w_1 W_{T_1 \cap \cdots \cap T_k} w_1^{-1}$$

for some $w_1 \in W$. Thus

$$\phi(W_{T_1 \cap \cdots \cap T_k}) \subset w'_1 W'_{T'_1 \cap \cdots \cap T'_k} {w'}_1^{-1} \subset w'_1 \phi(w_1) \phi(W_{T_1 \cap \cdots \cap T_k}) \phi(w_1^{-1}) {w'}_1^{-1}.$$

Hence $\phi(W_{T_1 \cap \cdots \cap T_k}) = w'_1 W'_{T'_1 \cap \cdots \cap T'_k} {w'}_1^{-1}$. Since $W_{T_1 \cap \cdots \cap T_k} \cong \mathbb{Z}_2^{|T_1 \cap \cdots \cap T_k|}$ and $W'_{T'_1 \cap \cdots \cap T'_k} \cong \mathbb{Z}_2^{|T'_1 \cap \cdots \cap T'_k|}$, we obtain that $|T_1 \cap \cdots \cap T_k| = |T'_1 \cap \cdots \cap T'_k|$. $\square$

Proof of Theorem 1.2. Let $T_1, \dots, T_n$ be the maximal spherical subsets of S and let $T'_1, \dots, T'_n$ be the maximal spherical subsets of S' such that $\phi(W_{T_i}) \sim W'_{T'_i}$ for each i. By (0) and (1), either $W_{T_i} \cong \mathbb{Z}_2^{|T_i|}$, or $T_i = \{s_i, t_i\}$ and $m(s_i, t_i)$ is odd. By Lemma 2.6, Lemma 3.2, (2), (3), ($2'$) and ($3'$), there exists a bijection $\psi : S \to S'$ such that

$$\psi(\bigcap_{i \in I} T_i) = \bigcap_{i \in I} T'_i$$

for each $I \subset \{1, \dots, n\}$. Then the bijection $\psi : S \to S'$ induces an isomorphism between (W,S) and (W',S'). $\square$

4. REMARK

The following theorem has been proved in [4].

Theorem 4.1 ([4]). *Every Coxeter system* (W, S) *which satisfies the conditions* (1), (2) *and* (3) *in Theorem 1.2 is reflection rigid.*

Radcliffe has proved the following theorem in [6].

Theorem 4.2 ([6]). *Let* (W, S) *be a Coxeter system such that* $m(s,t) \in \{2\} \cup 4\mathbb{N}$ *for each* $s \neq t \in S$*. Then the Coxeter group* W *is rigid.*

Here the following problem arises as an extension of Theorems 1.2 and 4.2.

Problem. Let (W, S) be a Coxeter system. Suppose that

- ($\bar{0}$) for each $s, t \in S$ such that $m(s,t)$ is even, $m(s,t) \in \{2\} \cup 4\mathbb{N}$,
- (1) for each $s \neq t \in S$ such that $m(s,t)$ is odd, $\{s,t\}$ is a maximal spherical subset of S,
- (2) there does not exist a three-points subset $\{s,t,u\} \subset S$ such that $m(s,t)$ and $m(t,u)$ are odd, and
- (3) for each $s \neq t \in S$ such that $m(s,t)$ is odd, the number of maximal spherical subsets of S intersecting with $\{s,t\}$ is at most two.

Is it the case that the Coxeter group W is rigid?

REFERENCES

[1] N. Bourbaki, *Groupes et Algebrès de Lie*, Chapters IV-VI, Masson, Paris, 1981.

[2] N. Brady, J.P. McCammond, B. Mühlherr and W.D. Neumann, *Rigidity of Coxeter groups and Artin groups*, Geom. Dedicata **94** (2002), 91–109.

[3] R. Charney and M.W. Davis, *When is a Coxeter system determined by its Coxeter group?* J. London Math. Soc. **61 (no.2)** (2000), 441–461.

[4] T. Hosaka, *A class of reflection rigid Coxeter systems*, preprint.

[5] D. Radcliffe, *Rigidity of right-angled Coxeter groups*, preprint.

[6] ———, *Unique presentation of Coxeter groups and related groups*, Ph.D. thesis, The University of Wisconsin-Milwaukee, 2001.

DEPARTMENT OF MATHEMATICS, UTSUNOMIYA UNIVERSITY, UTSUNOMIYA, 321-8505, JAPAN

E-mail address: `hosaka@cc.utsunomiya-u.ac.jp`